

DZD Publikationen 2010

A simplified method to assess affinity of insulin autoantibodies.

Achenbach P, Guo LH, Gick C, Adler K, Krause S, Bonifacio E, Colman PG, Ziegler AG. Clin Immunol. (2010);137:415. ([Abstract PubMed](#))

Improved insulin sensitivity, preserved beta cell function and improved whole-body glucose metabolism after low-dose growth hormone replacement therapy in adults with severe growth hormone deficiency: a pilot study.

Arafat AM, Möhlig M, Weickert MO, Schöfl C, Spranger J, Pfeiffer AF. Diabetologia. (2010) ;53:1304 ([Artikel](#)) ([pdf](#))

Monocyte gene-expression profiles associated with childhood-onset type 1 diabetes and disease risk: a study of identical twins.

Beyan H, Drexhage RC, van der Heul NL, de WH, Padmos RC, Schloot NC, Drexhage HA, Leslie RD. (2010) Diabetes 59: 1751. ([Artikel](#)) ([pdf](#))

TOSS - Multiple hypothesis testing in an open software system.

Blanchard G, Dickhaus T, Hack N, Konietzschke F, Rohmeyer K, Rosenblatt J, Scheer M, Werft W. (2010) Workshop and Conference Proceedings Journal of Machine Learning Research 11: 12. ([pdf](#))

Probiotics and obesity.

Blaut M, Bischoff SC. Ann Nutr Metab. (2010);57 Suppl:20 ([Abstract PubMed](#))

Harmonization of glutamic acid decarboxylase and islet antigen-2 autoantibody assays for national institute of diabetes and digestive and kidney diseases consortia.

Bonifacio E, Yu L, Williams AK, Eisenbarth GS, Bingley PJ, Marcovina SM, Adler K, Ziegler AG, Mueller PW, Schatz DA, Krischer JP, Steffes MW, Akolkar B. J Clin Endocrinol Metab. (2010) ;95:3360. ([Artikel](#)) ([pdf](#))

Advances in the prediction and natural history of type 1 diabetes.

Bonifacio E, Ziegler AG. Endocrinol Metab Clin North Am. (2010);39:513 ([Abstract PubMed](#))

Acute elevation of plasma lipids does not affect ATP synthesis in human skeletal muscle.

Brehm A, Krssak M, Schmid AI, Nowotny P, Waldhausl W, Roden M. (2010) Am. J. Physiol. Endocrinol. Metab. 299: E33. ([Abstract PubMed](#))

Insel(zell)-Transplantationen bei Typ-1-Diabetes-Patienten

Brendel MD, Ludwig B, Eckhard M, Bretzel RG. Transplantationen bei Typ-1-Diabetikern: Diabetologe (2010);6:357

COUP-TFII is regulated by high glucose in endothelial cells.

Brunssen C, Korten S, Brux M, Seifert S, Roesler J, Bornstein SR, Morawietz H, Goettsch W. Horm Metab Res. (2010) ;42:81. ([Abstract PubMed](#))

Birth order and childhood type 1 diabetes: a pooled analysis of 31 observational studies.

Cardwell CR, Stene LC, Joner G, Bulsara MK, Cinek O, Rosenbauer J, Ludvigsson J, Svensson J, Goldacre MJ, Waldhoer T, Jarosz-Chobot P, Gimeno SG, Chuang LM, Roberts CL, Parslow RC, Wadsworth EJ, Chetwynd A, Brigis G, Urbonaite B, Sipetic S, Schober E, Devoti G, Ionescu-Tirgoviste C, de Beaufort CE, Stoyanov D, Buschard K, Radon K, Glatthaar C, Patterson CC. (2010) Int.J.Epidemiol. [\(Abstract PubMed\)](#)

Maternal age at birth and childhood type 1 diabetes: a pooled analysis of 30 observational studies.

Cardwell CR, Stene LC, Joner G, Bulsara MK, Cinek O, Rosenbauer J, Ludvigsson J, Jane M, Svensson J, Goldacre MJ, Waldhoer T, Jarosz-Chobot P, Gimeno SGA, Chuang LM, Parslow RC, Wadsworth EJK, Chetwynd A, Pozzilli P, Brigis G, Urbonaite B, Sipetic S, Schober E, Devoti G, Ionescu-Tirgoviste C, de Beaufort CE, Stoyanov D, Buschard K, Patterson CC. (2010) Diabetes 59: 486. [\(Artikel\)](#) [\(pdf\)](#)

Birthweight and the risk of childhood-onset type 1 diabetes: a meta-analysis of observational studies using individual patient data.

Cardwell CR, Stene LC, Joner G, Davis EA, Cinek O, Rosenbauer J, Ludvigsson J, Castell C, Svensson J, Goldacre MJ, Waldhoer T, Polanska J, Gimeno SGA, Chuang LM, Parslow RC, Wadsworth EJK, Chetwynd A, Pozzilli P, Brigis G, Urbonaite B, Sipetic S, Schober E, Ionescu-Tirgoviste C, de Beaufort CE, Stoyanov D, Buschard K, Patterson CC. (2010) Diabetologia 53: 641. [\(Abstract PubMed\)](#)

Accelerated increase in serum interleukin-1 receptor antagonist starts 6 years before diagnosis of type 2 diabetes: Whitehall II prospective cohort study.

Carstensen M, Herder C, Kivimaki M, Jokela M, Roden M, Shipley MJ, Witte DR, Brunner EJ, Tabak AG. (2010) Diabetes 59: 1222. [\(Artikel\)](#)

Macrophage inhibitory cytokine-1 is increased in individuals before type 2 diabetes diagnosis but is not an independent predictor of type 2 diabetes: the whitehall II study.

Carstensen M, Herder C, Brunner EJ, Strassburger K, Tabak AG, Roden M, Witte DR. (2010) Eur.J.Endocrinol. 162: 913. [\(Abstract PubMed\)](#)

Within-trial economic evaluation of diabetes-specific cognitive behaviour therapy in patients with type 2 diabetes and subthreshold depression.

Chernyak C, Kulzer B, Hermanns N, Schmitt A, Gahr A, Haak T, Kruse J, Ohmann C, Scheer M, Giani G, Icks A. (2010) BMC.Public Health 10: 625. [\(Artikel\)](#) [\(pdf\)](#)

PPARgamma-dependent regulation of adenylate cyclase 6 amplifies the stimulatory effect of cAMP on renin gene expression.

Desch M, Schubert T, Schreiber A, Mayer S, Friedrich B, Artunc F, Todorov VT. Mol Endocrinol. (2010);24:2139 [\(Artikel\)](#) [\(pdf\)](#)

Metabolic control in children with diabetes mellitus type 1.

Dost A, Rosenbauer J, Rohrer T, Knerr I, Gerstl EM, Holl RW, for the Initiative DPV Science and the Competence Network Diabetes. (2010) Eur.J.Endocrinol. 6: 35.

Diet-induced gene expression of isolated pancreatic islets from a polygenic mouse model of the metabolic syndrome.

Dreja T, Jovanovic Z, Rasche A, Kluge R, Herwig R, Tung YC, Joost HG, Yeo GS, Al-Hasani H. Diabetologia. (2010);53:3 ([Artikel](#)) ([pdf](#))

New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk.

Dupuis, J. et al. (2010) Nature Genetics 42: 105. ([Artikel](#)) ([pdf](#))

Posttransplantationsdiabetes

Eckhard M, Dieterle C, Brendel MD, Bretzel RG, Landgraf R. Diabetologe (2010) 6:460

Impaired insulin turnover in islets from type 2 diabetic patients.

Eehalt F, Knoch K, Erdmann K, Krautz C, Jäger M, Steffen A, Wegbrod C, Meisterfeld R, Kersting S, Bergert H, Kuhlisch E, Bornstein S, Bonifacio E, Saeger HD, Solimena M. Islets. (2010) ;2:30. ([pdf](#))

Pigment epithelium-derived factor is one of the most abundant proteins secreted by human adipocytes and induces insulin resistance and inflammatory signaling in muscle and fat cells.

Famulla S, Lamers D, Hartwig S, Passlack W, Horrighs A, Cramer A, Lehr S, Sell H, Eckel J. (2010) Int.J.Obesity (in press). ([PubMed Abstract](#))

Absence of intestinal microbiota does not protect mice from diet-induced obesity.

Fleissner CK, Huebel N, Abd El-Bary MM, Loh G, Klaus S, Blaut M. Br J Nutr. (2010);104:919 ([Abstract PubMed](#))

Processing of food pictures: influence of hunger, gender and calorie content.

Frank S, Laharnar N, Kullmann S, Veit R, Canova C, Hegner YL, Fritsche A, Preissl H. Brain Res. (2010)2;1350:159 ([Artikel](#))

Predictors of diabetic ketoacidosis in children and adolescents with type 1 diabetes.

Fritsch MJ, Rosenbauer J, Schober E, Neu A, Placzek K, Holl RW. (2010) Experience from a large multicenter data base. Journal of Pediatrics: in press.

IL-6 deficiency in mice neither impairs induction of metabolic genes in the liver nor affects blood glucose levels during fasting and moderately intense exercise.

Fritsche L, Hoene M, Lehmann R, Ellingsgaard H, Hennige AM, Pohl AK, Häring HU, Schleicher ED, Weigert C. Diabetologia. (2010);53:1732 ([Artikel](#)) ([pdf](#))

Comparison between a basal-bolus and a premixed insulin regimen in individuals with type 2 diabetes-results of the GINGER study.

Fritsche A, Larbig M, Owens D, Häring HU; GINGER study group. Diabetes Obes Metab. (2010);12:115 ([Artikel](#)) ([pdf](#))

Mouse Phenotyping

Fuchs et al. Methods (2010);12

Benefit-risk assessment of exenatide in the therapy of type 2 diabetes mellitus.

Gallwitz B. Drug Saf. (2010);33:87 ([Abstract PubMed](#))

The evolving place of incretin-based therapies in type 2 diabetes.

Gallwitz B. Pediatr Nephrol. (2010);25:1207 ([Artikel](#)) ([pdf](#))

Future perspectives for insulinotropic agents in the treatment of type 2 diabetes-DPP-4 inhibitors and sulphonylureas.

Gallwitz B, Häring HU. Diabetes Obes Metab. (2010);12:1 ([Artikel](#)) ([pdf](#))

Changes in body composition after 9 months of treatment with exenatide twice daily versus glimepiride: comment letter on Jendle et al.

Gallwitz B, Haupt A, Kraus P, Peters N, Petto H, Dotta F, Petto H, Poll L, Rose L, Schernthaner G. Diabetes Obes Metab. (2010);12:1127 ([Abstract PubMed](#))

Adding liraglutide to oral antidiabetic drug therapy: onset of treatment effects over time.

Gallwitz B, Vaag A, Falahati A, Madsbad S. Int J Clin Pract. (2010);64:267 ([Artikel](#)) ([pdf](#))

[Incretins: do they exert cardiovascular effects?].

[Article in German]

Gallwitz B. Herz. (2010);35:130 ([pdf](#))

Blood glucose testing and primary prevention of diabetes mellitus type 2--evaluation of the effect of evidence based patient information.

Genz J, Haastert B, Meyer G, Steckelberg A, Muller H, Verheyen F, Cole D, Rathmann W, Nowotny B, Roden M, Giani G, Mielck A, Ohmann C, Icks A. (2010) BMC.Public Health 10: 15. ([Artikel](#)) ([pdf](#))

Reduced incidence of blindness in relation to diabetes mellitus in southern Germany?

Genz J, Scheer M, Trautner C, Zollner I, Giani G, Icks A. (2010) Diabet.Med. 27: 1138. ([Abstract PubMed](#))

Saxagliptin is non-inferior to glipizide in patients with type 2 diabetes mellitus inadequately controlled on metformin alone: a 52-week randomised controlled trial.

Göke B, Gallwitz B, Eriksson J, Hellqvist A, Gause-Nilsson I; D1680C00001 Investigators. Int J Clin Pract. (2010);64:1619 ([Abstract PubMed](#))

Type 2 Diabetes in Octogenarians Is Associated with Decreased Low Molecular Weight Adiponectin.

Graessler J, Gruber M, Radke RB, Kopprasch S, Schwarz PE, Kamke W, Bornstein SR, Fischer S. Gerontology. (2010) 11. [Epub ahead of print] ([Abstract PubMed](#))

Association of genetic variation in KCNQ1 with type 2 diabetes in the KORA surveys.

Grallert H, Herder C, Marzi C, Meisinger C, Wichmann HE, Rathmann W, Illig T. (2010) Horm.Metab Res. 42: 149. ([Abstract PubMed](#))

Insulin modulates food-related activity in the central nervous system.

Guthoff M, Grichisch Y, Canova C, Tschritter O, Veit R, Hallschmid M, Häring HU, Preissl H, Hennige AM, Fritsche A. J Clin Endocrinol Metab. (2010);95:748 ([Artikel](#)) ([pdf](#))

A polymorphism in the gene encoding AdipoR1 affects olfactory recognition.

Guthoff M, Tschritter O, Berg D, Liepelt I, Schulte C, Machicao F, Breer H, Haering HU, Fritsche A. Int J Obes (Lond). (2010) ([Artikel](#)) ([pdf](#))

Gene variants of TCF7L2 influence weight loss and body composition during lifestyle intervention in a population at risk for type 2 diabetes.

Haupt A, Thamer C, Heni M, Ketterer C, Machann J, Schick F, Machicao F, Stefan N, Claussen CD, Häring HU, Fritsche A, Staiger H. Diabetes. (2010);59:747 ([Artikel](#)) ([pdf](#))

Novel obesity risk loci do not determine distribution of body fat depots: a whole-body MRI/MRS study.

Haupt A, Thamer C, Heni M, Machicao F, Machann J, Schick F, Stefan N, Fritsche A, Häring HU, Staiger H. Obesity (Silver Spring). (2010);18:1212 ([Artikel](#)) ([pdf](#))

Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution.

Heid et al. Nat Genet. (2010) ;42:949. ([Artikel](#)) ([pdf](#))

Association of obesity risk SNPs in PCSK1 with insulin sensitivity and proinsulin conversion.

Heni M, Haupt A, Schäfer SA, Ketterer C, Thamer C, Machicao F, Stefan N, Staiger H, Häring HU, Fritsche A. (2010) BMC Med Genet. 9: 11. ([Artikel](#)) ([pdf](#))

The impact of genetic variation in the G6PC2 gene on insulin secretion depends on glycemia.

Heni M, Ketterer C, Hart LM, Ranta F, van Haeften TW, Eekhoff EM, Dekker JM, Boomsma DI, Nijpels G, Kramer MH, Diamant M, Simonis-Bik AM, Heine RJ, de Geus EJ, Schäfer SA, Machicao F, Ullrich S, Thamer C, Stefan N, Staiger H, Häring HU, Fritsche A. (2010) J Clin Endocrinol Metab. 95: E479. ([Abstract PubMed](#))

Glycemia determines the effect of type 2 diabetes risk genes on insulin secretion.

Heni M, Ketterer C, Thamer C, Herzberg-Schäfer SA, Guthoff M, Stefan N, Machicao F, Staiger H, Fritsche A, Häring HU. (2010) Diabetes 59: 3247. ([Abstract PubMed](#))

Pancreatic fat is negatively associated with insulin secretion in individuals with impaired fasting glucose and/or impaired glucose tolerance: a nuclear magnetic resonance study.

Heni M, Machann J, Staiger H, Schwenzer NF, Peter A, Schick F, Claussen CD, Stefan N, Häring HU, Fritsche A. Diabetes Metab Res Rev. (2010);26:200 ([Abstract PubMed](#))

Variants in the CD36 Gene Locus Determine Whole-Body Adiposity, but Have No Independent Effect on Insulin Sensitivity.

Heni M, Müssig K, Machicao F, Machann J, Schick F, Claussen CD, Stefan N, Fritsche A, Häring HU, Staiger H. Obesity (Silver Spring). (2010) 21. [Epub ahead of print] ([Artikel](#)) ([pdf](#))

Enforced expression of protein kinase C in skeletal muscle causes physical inactivity, fatty liver and insulin resistance in the brain.

Hennige AM, Heni M, Machann J, Staiger H, Sartorius T, Hoene M, Lehmann R, Weigert C, Peter A, Bornemann A, Kroeber S, Pujol A, Franckhauser S, Bosch F, Schick F, Lammers R, Häring HU. *J Cell Mol Med.* (2010);14:903 ([Abstract PubMed](#))

Overexpression of kinase-negative protein kinase Cdelta in pancreatic beta-cells protects mice from diet-induced glucose intolerance and beta-cell dysfunction.

Hennige AM, Ranta F, Heinzemann I, Düfer M, Michael D, Braumüller H, Lutz SZ, Lammers R, Drews G, Bosch F, Häring HU, Ullrich S. *Diabetes.* (2010);59:119 ([Artikel](#)) ([pdf](#))

Evaluation of fasting state-/oral glucose tolerance test-derived measures of insulin release for the detection of genetically impaired β -cell function.

Herzberg-Schäfer SA, Staiger H, Heni M, Ketterer C, Guthoff M, Kantartzis K, Machicao F, Stefan N, Häring HU, Fritsche A. (2010) *PLoS One* 5: e14194. ([Artikel](#)) ([pdf](#))

Altered GLUT4 trafficking in adipocytes in the absence of the GTPase Arfrp1.

Hesse D, Hommel A, Jaschke A, Moser M, Bernhardt U, Zahn C, Kluge R, Wittschen P, Gruber AD, Al-Hasani H, Joost HG, Schürmann A. (2010) *Biochem Biophys Res Commun.* 394: 896. ([Abstract PubMed](#))

Decision trees as a simple-to-use and reliable tool to identify individuals with impaired glucose metabolism or type 2 diabetes mellitus.

Hische M, Luis-Dominguez O, Pfeiffer AF, Schwarz PE, Selbig J, Spranger J. *Eur J Endocrinol.* (2010) ;163:565. ([Abstract](#))

The effect of diabetes on incidence and mortality in end-stage renal disease in Germany.

Hoffmann F, Haastert B, Koch M, Giani G, Icks A. (2010) *Nephrol.Dial.Transplant.* (in press). ([Abstract PubMed](#))

The ARF-like GTPase ARFRP1 is essential for lipid droplet growth and is involved in the regulation of lipolysis.

Hommel A, Hesse D, Völker W, Jaschke A, Moser M, Engel T, Blüher M, Zahn C, Chadt A, Ruschke K, Vogel H, Kluge R, Robenek H, Joost HG, Schürmann A. (2010) *Mol Cell Biol.* 30: 1231. ([Artikel](#)) ([pdf](#))

Activation of the mitogen-activated protein kinase (MAPK) signalling pathway in the liver of mice is related to plasma glucose levels after acute exercise.

Hoene M, Franken H, Fritsche L, Lehmann R, Pohl AK, Häring HU, Zell A, Schleicher ED, Weigert C. *Diabetologia.* (2010) ;53:1131 ([Artikel](#)) ([pdf](#))

The stress response of the liver to physical exercise.

Hoene M, Weigert C. *Exerc Immunol Rev.* (2010);16:163 ([pdf](#))

Meta-analysis of individual patient data to assess the risk of hypoglycaemia in people with type 2 diabetes using NPH insulin or insulin glargine.

Home PD, Fritsche A, Schinzel S, Massi-Benedetti M. Diabetes Obes Metab. (2010);12:772 ([Abstract PubMed](#))

Lipidomics analysis reveals efficient storage of hepatic triacylglycerides enriched in unsaturated fatty acids after one bout of exercise in mice.

Hu C, Hoene M, Zhao X, Häring HU, Schleicher E, Lehmann R, Han X, Xu G, Weigert C. (2010) PLoS One 5: e13318. ([Artikel](#)) ([pdf](#))

Stroke in the diabetic and non-diabetic population in Germany: relative and attributable risks, 2005-2007.

Icks A, Scheer M, Genz J, Giani G, Glaeske G, Hoffmann F. (2010) J.Diabetes Complications (in press). ([Abstract PubMed](#))

Costs of dialysis--a regional population-based analysis.

Icks A, Haastert B, Gandjour A, Chernyak N, Rathmann W, Giani G, Rump LC, Trapp R, Koch M. (2010) Nephrol.Dial.Transplant. 25: 1647. ([Abstract PubMed](#))

Detailed physiologic characterization reveals diverse mechanisms for novel genetic Loci regulating glucose and insulin metabolism in humans.

Ingelsson E, Langenberg C, Hivert MF, Prokopenko I, Lyssenko V, Dupuis J, Mägi R, Sharp S, Jackson AU, Assimes TL, Shrader P, Knowles JW, Zethelius B, Abbasi FA, Bergman RN, Bergmann A, Berne C, Boehnke M, Bonnycastle LL, Bornstein SR, Buchanan TA, Bumpstead SJ, Böttcher Y, Chines P, Collins FS, Cooper CC, Dennison EM, Erdos MR, Ferrannini E, Fox CS, Graessler J, Hao K, Isomaa B, Jameson KA, Kovacs P, Kuusisto J, Laakso M, Ladenvall C, Mohlke KL, Morken MA, Narisu N, Nathan DM, Pascoe L, Payne F, Petrie JR, Sayer AA, Schwarz PE, Scott LJ, Stringham HM, Stumvoll M, Swift AJ, Syvänen AC, Tuomi T, Tuomilehto J, Tönjes A, Valle TT, Williams GH, Lind L, Barroso I, Quertermous T, Walker M, Wareham NJ, Meigs JB, McCarthy MI, Groop L, Watanabe RM, Florez JC; MAGIC investigators. Diabetes. (2010);59:1266 ([Artikel](#)) ([pdf](#))

A genome-wide perspective of genetic variation in human metabolism.

Illig T, Gieger C, Zhai G, Römisch-Margl W, Wang-Sattler R, Prehn C, Altmaier E, Kastenmüller G, Kato BS, Mewes HW, Meitinger T, de Angelis MH, Kronenberg F, Soranzo N, Wichmann HE, Spector TD, Adamski J, Suhre K. (2010) Nature Genetics 42: 137. ([Abstract PubMed](#))

Effects of long-term soluble vs. insoluble dietary fiber intake on high-fat diet-induced obesity in C57BL/6J mice.

Isken F, Klaus S, Osterhoff M, Pfeiffer AF, Weickert MO. J Nutr Biochem. (2010);21:278 ([Artikel](#))

Impairment of fat oxidation under high- vs. low-glycemic index diet occurs before the development of an obese phenotype.

Isken F, Klaus S, Petzke KJ, Loddenkemper C, Pfeiffer AF, Weickert MO. Am J Physiol Endocrinol Metab. (2010);298:E287 ([Artikel](#)) ([pdf](#))

Recent economic findings on childhood obesity: cost-of-illness and cost-effectiveness of interventions.

John J, Wenig CM, Wolfenstetter SB. *Curr Opin Clin Nutr Metab Care* (2010) 13: 305. ([Abstract PubMed](#))

The genetic basis of obesity and type 2 diabetes: lessons from the new zealand obese mouse, a polygenic model of the metabolic syndrome.

Joost HG. *Results Probl Cell Differ.* (2010);52:1 ([Abstract PubMed](#))

Diabetes Mellitus Typ 2: Risikobestimmung wird präzisiert.

Joost HG, Fritsche A, Häring HU, Pfeiffer AFH, Roden M, Schulze MB. (2010) *Deutsches Ärzteblatt* 107: A 600-2. ([Artikel](#)) ([pdf](#))

Association of adiponectin, interleukin (IL)-1ra, inducible protein 10, IL-6 and number of islet autoantibodies with progression patterns of type 1 diabetes the first year after diagnosis.

Kaas A, Pflieger C, Hansen L, Buschard K, Schloot NC, Roep BO, Mortensen HB. (2010) *Clin Exp Immunol.* 161: 444. ([Abstract PubMed](#))

Impaired insulin stimulation of muscular ATP production in patients with type 1 diabetes.

Kaceroovsky M, Brehm A, Chmelik M, Schmid AI, Szendroedi J, Kaceroovsky-Bielez G, Nowotny P, Lettner A, Wolzt M, Jones JG, Roden M. *J Intern Med.* (2011);269:189. ([Abstract PubMed](#))

The impact of liver fat vs visceral fat in determining categories of prediabetes.

Kantartzis K, Machann J, Schick F, Fritsche A, Häring HU, Stefan N. *Diabetologia.* (2010);53:882 ([Artikel](#)) ([pdf](#))

Environmental and genetic determinants of fatty liver in humans.

Kantartzis K, Schick F, Häring HU, Stefan N. *Dig Dis.* (2010);28:169 ([Abstract PubMed](#))

Leptin, adiponectin, their ratio and risk of Coronary Heart Disease: results from the MONICA/KORA Augsburg Study 1984-2002.

Karakas M, Zierer A, Herder C, Baumert J, Meisinger C, Koenig W, Thorand B. (2010) *Atherosclerosis* 209: 220. ([Abstract PubMed](#))

Long-acting insulin analogs and the risk of diabetic ketoacidosis in children and adolescents with type 1 diabetes. A prospective study of 10,682 patients from 271 institutions.

Karges B, Kapellen T, Neu A, Hofer SE, Rohrer T, Rosenbauer J, Wolf J, Holl RW. (2010) *Diabetes Care* 33: 1031. ([Artikel](#)) ([pdf](#))

Effects of coffee consumption on subclinical inflammation and other risk factors for type 2 diabetes: a clinical trial.

Kempf K, Herder C, Erlund I, Kolb H, Martin S, Carstensen M, Koenig W, Sundvall J, Bidel S, Kuha S, Tuomilehto J. (2010) *Am.J.Clin.Nutr.* 91: 950. ([Abstract PubMed](#))

Vaspin (SERPINA12) genotypes and risk of type 2 diabetes: Results from the MONICA/KORA studies.

Kempf K, Rose B, Illig T, Rathmann W, Strassburger K, Thorand B, Meisinger C, Wichmann HE, Herder C, Vollmert C. (2010) Exp.Clin.Endocrinol.Diabetes 118: 184. ([Abstract PubMed](#))

European lactase persistence genotype shows evidence of association with increase in body mass index.

Kettunen J, Silander K, Saarela O, Amin N, Müller M, Timpson N, Surakka I, Ripatti S, Laitinen J, Hartikainen AL, Pouta A, Lahermo P, Anttila V, Männistö S, Jula A, Virtamo J, Salomaa V, Lehtimäki T, Raitakari O, Gieger C, Wichmann EH, Van Duijn CM, Smith GD, McCarthy MI, Järvelin MR, Perola M, Peltonen L.

Hum Mol Genet. (2010) 19: 1129. ([Artikel](#)) ([pdf](#))

Acute, short-term hyperinsulinemia increases olfactory threshold in healthy subjects.

Ketterer C, Heni M, Thamer C, Herzberg-Schäfer SA, Häring HU, Fritsche A. Int J Obes (Lond). (2010) Nov 23. [Epub ahead of print] doi: 10.1038 ([Abstract PubMed](#))

Genetic variation within the NR1H2 gene encoding liver X receptor β associates with insulin secretion in subjects at increased risk for type 2 diabetes.

Ketterer C, Müssig K, Machicao F, Stefan N, Fritsche A, Häring HU, Staiger H.J Mol Med. (2011) ;89:75. ([Artikel](#)) ([pdf](#))

[Insulin pump therapy: indications and risks - Case 7/2010].

[Article in German]

Ketterer C, Müssig K, Gerstenecker AK, Fritsche A. Dtsch Med Wochenschr. (2010);135 ([Abstract PubMed](#))

The rs1800629 polymorphism in the TNF gene interacts with physical activity on the changes in C-reactive protein levels in the finnish diabetes prevention study.

Kilpelainen TO, Laaksonen DE, Lakka TA, Herder C, Koenig W, Lindstrom J, Eriksson JG, Uusitupa M, Kolb H, Laakso M, Tuomilehto J. (2010) Exp Clin Endocrinol Diabetes 118: 757. ([Abstract PubMed](#))

Therapieoptionen bei Hypophysenadenomen

Knappe U, Brendel MD. MedWelt (2010);61:7

Phosphorylation of sterol regulatory element-binding protein (SREBP)-1a links growth hormone action to lipid metabolism in hepatocytes.

Kotzka J, Knebel B, Avci H, Jacob S, Nitzgen U, Jockenhovel F, Heeren J, Haas J, Muller-Wieland D. (2010) Atherosclerosis 213: 156. ([Abstract PubMed](#))

Socioeconomic status is not associated with type 2 diabetes incidence in an elderly population in Germany: KORA S4/F4 Cohort Study.

Kowall B, Rathmann W, Strassburger K, Meisinger C, Holle R, Mielck A. (2010) J.Epidemiol.Community Health 25: 393. ([Artikel](#)) ([pdf](#))

Association of passive and active smoking with incident type 2 diabetes mellitus in the elderly population: the KORA S4/F4 cohort study.

Kowall B, Rathmann W, Strassburger K, Heier M, Holle R, Thorand B, Giani G, Peters A, Meisinger C. (2010) Eur.J.Epidemiol. 25: 393. ([Abstract PubMed](#))

Effect of sedentary behaviour and vigorous physical activity on segment-specific carotid wall thickness and its progression in a healthy population.

Kozakova M, Palombo C, Morizzo C, Nolan JJ, Konrad T, Balkau B, and the RISC Investigators (Roden M). (2010) Eur.Heart J. 31: 1511. ([Abstract PubMed](#))

Glucokinase-activating GCKR polymorphisms increase plasma levels of triglycerides and free fatty acids, but do not elevate cardiovascular risk in the Ludwigshafen Risk and Cardiovascular Health Study.

Kozian DH, Barthel A, Cousin E, Brunnhöfer R, Anderka O, März W, Böhm B, Winkelmann B, Bornstein SR, Schmol D. Horm Metab Res. (2010) ;42:502 ([Abstract PubMed](#))

Traffic-related air pollution and incident type 2 diabetes: results from the SALIA cohort study.

Krämer U, Herder C, Sugiri D, Strassburger K, Schikowski T, Ranft U, Rathmann W. (2010) Environ.Health Perspect. 118: 1273. ([Artikel](#)) ([pdf](#))

Zinc transporter 8 antibodies complement GAD and IA-2 antibodies in the identification and characterization of adult-onset autoimmune diabetes: Non Insulin Requiring Autoimmune Diabetes (NIRAD) 4.

Lampasona V, Petrone A, Tiberti C, Capizzi M, Spoletini M, di Pietro S, Songini M, Bonicchio S, Giorgino F, Bonifacio E, Bosi E, Buzzetti R; Non Insulin Requiring Autoimmune Diabetes (NIRAD) Study Group. Diabetes Care. (2010) ;33:104 ([Artikel](#)) ([pdf](#))

Complement-mediated inhibition of neovascularization reveals a point of convergence between innate immunity and angiogenesis.

Langer HF, Chung KJ, Orlova VV, Choi EY, Kaul S, Kruhlak MJ, Alatsatianos M, Deangelis RA, Roche PA, Magotti P, Li X, Economopoulou M, Rafail S, Lambiris JD, Chavakis T. Blood. (2010)25;116:4395 ([Artikel](#)) ([pdf](#))

Do FADS genotypes enhance our knowledge about fatty acid related phenotypes?

Lattka E, Illig T, Heinrich J, Koletzko B.Clin Nutr.(2010) 29: 277. ([Abstract PubMed](#))

Sex steroids affect triglyceride handling, glucose-dependent insulinotropic polypeptide, and insulin sensitivity: a 1-week randomized clinical trial in healthy young men.

Lapauw B, Ouwens M, 't Hart LM, Wuyts B, Holst JJ, T'Sjoen G, Kaufman JM, Ruige JB. (2010) Diabetes Care 33: 1831. ([Artikel](#)) ([pdf](#))

Association of the FTO gene variant (rs9939609) with cardiovascular disease in men with abnormal glucose metabolism - The Finnish Diabetes Prevention Study.

Lappalainen T, Kolehmainen M, Schwab US, Tolppanen AM, Stancakova A, Lindstrom J, Eriksson JG, Keinanen-Kiukaanniemi S, Aunola S, Ilanne-Parikka P, Herder C, Koenig W, Gylling H, Kolb H, Tuomilehto J, Kuusisto J, Uusitupa M. (2010) Nutr.Metab Cardiovasc.Dis. Epub Apr 2010. ([Abstract PubMed](#))

Intra-epidermal nerve fibre density at the distal leg: A worldwide normative reference study.

Lauria G, Bakkers M, Schmitz C, Lombardi R, Penza P, Devigili G, Smith A, Hsieh S-T, Mellgren S, Umapathi T, Ziegler D, Faber C, Merckies I. (2010) J Peripher Nerv Syst 15: 202. ([Abstract PubMed](#))

Medium chain acylcarnitines dominate the metabolite pattern in humans under moderate intensity exercise and support lipid oxidation.

Lehmann R, Zhao X, Weigert C, Simon P, Fehrenbach E, Fritsche J, Machann J, Schick F, Wang J, Hoene M, Schleicher ED, Häring HU, Xu G, Niess AM. (2010) PLoS One. (7): e11519. ([Artikel](#)) ([pdf](#))

Take action to prevent diabetes--the IMAGE toolkit for the prevention of type 2 diabetes in Europe.

Lindström J. et al. Horm Metab Res. (2010) ;42 Suppl 1:S37 ([Abstract PubMed](#))

Tamoxifen-independent recombination in the RIP-CreER mouse.

Liu Y, Suckale J, Masjkur J, Magro MG, Steffen A, Anastassiadis K, Solimena M. PLoS One. (2010) 22;5:e13533 ([Artikel](#)) ([pdf](#))

Insulin sensitivity is reflected by characteristic metabolic fingerprints--a Fourier transform mass spectrometric non-targeted metabolomics approach.

Lucio M, Fekete A, Weigert C, Wägele B, Zhao X, Chen J, Fritsche A, Häring HU, Schleicher ED, Xu G, Schmitt-Kopplin P, Lehmann R. (2010) PLoS One 5: e13317. ([Artikel](#)) ([pdf](#))

Agonist of growth hormone-releasing hormone as a potential effector for survival and proliferation of pancreatic islets.

Ludwig B, Ziegler CG, Schally AV, Richter C, Steffen A, Jabs N, Funk RH, Brendel MD, Block NL, Ehrhart-Bornstein M, Bornstein SR. (2010) Proc Natl Acad Sci U S A. 107: 12623. ([Artikel](#)) ([pdf](#))

A novel device for islet transplantation providing immune protection and oxygen supply.

Ludwig B, Zimmerman B, Steffen A, Yavriants K, Azarov D, Reichel A, Vardi P, German T, Shabtay N, Rotem A, Evron Y, Neufeld T, Mimon S, Ludwig S, Brendel MD, Bornstein SR, Barkai U. (2010) Horm Metab Res. 42: 918. ([Abstract PubMed](#))

Islet versus pancreas transplantation in type 1 diabetes: competitive or complementary?

Ludwig B, Ludwig S, Steffen A, Saeger HD, Bornstein SR. Curr Diab Rep. (2010) ;10:506. ([Artikel](#)) ([pdf](#))

Follow-up whole-body assessment of adipose tissue compartments during a lifestyle intervention in a large cohort at increased risk for type 2 diabetes.

Machann J, Thamer C, Stefan N, Schwenzer NF, Kantartzis K, Häring HU, Claussen CD, Fritsche A, Schick F. Radiology. (2010);257:353 ([Abstract PubMed](#))

Isolation and preservation of peripheral blood mononuclear cells for analysis of islet antigen-reactive T cell responses: position statement of the T-Cell Workshop Committee of the Immunology of Diabetes Society.

Mallone R, Mannering SI, Brooks-Worrell BM, Durinovic-Belló I, Cilio CM, Wong FS, Schloot NC; T-Cell Workshop Committee, Immunology of Diabetes Society Clin Exp Immunol. (2011);163:33 ([Abstract PubMed](#))

Current approaches to measuring human islet-antigen specific T cell function in type 1 diabetes.

Mannering SI, Wong FS, Durinovic-Bello I, Brooks-Worrell B, Tree TI, Cilio CM, Schloot NC, Mallone R. (2010) Clin Exp Immunol. 162: 197. ([Abstract PubMed](#))

The use of statins potentiates the insulin-sensitizing effect of exercise training in obese males with and without Type 2 diabetes.

Meex RC, Phielix E, Schrauwen-Hinderling VB, Moonen-Kornips E, Schaart G, Schrauwen P, Hesselink MK. (2010) Clin.Sci.(Lond) 119: 293. ([Abstract PubMed](#))

Restoration of muscle mitochondrial function and metabolic flexibility in type 2 diabetes by exercise training is paralleled by increased myocellular fat storage and improved insulin sensitivity.

Meex RC, Schrauwen-Hinderling VB, Moonen-Kornips E, Schaart G, Mensink M, Phielix E, van de Weijer T, Sels JP, Schrauwen P, Hesselink MK. (2010) Diabetes 59: 572. ([Artikel](#)) ([pdf](#))

Prevalence of undiagnosed diabetes and impaired glucose regulation in 35-59-year-old individuals in Southern Germany: the KORA F4 Study.

Meisinger C, Strassburger K, Heier M, Thorand B, Baumeister SE, Giani G, Rathmann W. (2010) Diabet.Med. 27: 360. ([Abstract PubMed](#))

Co-graft of allogeneic immune regulatory neural stem cells (NPC) and pancreatic islets mediates tolerance, while inducing NPC-derived tumors in mice.

Melzi R, Antonioli B, Mercalli A, Battaglia M, Valle A, Pluchino S, Galli R, Sordi V, Bosi E, Martino G, Bonifacio E, Doglioni C, Piemonti L. PLoS One. (2010) 27;5:e10357. ([Artikel](#)) ([pdf](#))

Role of CCL2/MCP-1 in islet transplantation.

Melzi R, Mercalli A, Sordi V, Cantarelli E, Nano R, Maffi P, Sitia G, Guidotti LG, Secchi A, Bonifacio E, Piemonti L. Cell Transplant. (2010) ;19:1031 ([Abstract PubMed](#))

In situ synthesis of magnetic multiwalled carbon nanotube composites for the clean-up of (fluoro)quinolones from human plasma prior to ultrahigh pressure liquid chromatography analysis.

Morales-Cid G, Fekete A, Simonet BM, Lehmann R, Cárdenas S, Zhang X, Valcárcel M, Schmitt-Kopplin P. Anal Chem. (2010)1;82:2743 ([Artikel](#)) ([pdf](#))

The ENPP1 K121Q polymorphism determines individual susceptibility to the insulin-sensitising effect of lifestyle intervention.

Müssig K, Heni M, Thamer C, Kantartzis K, Machicao F, Stefan N, Fritsche A, Häring HU, Staiger H. Diabetologia. (2010);53:504 ([Artikel](#)) ([pdf](#))

Insulin signal transduction in normal cells and its role in carcinogenesis.

Müssig K, Häring HU. Exp Clin Endocrinol Diabetes. (2010);118:356 ([Abstract PubMed](#))

No association between variation in the NR4A1 gene locus and metabolic traits in white subjects at increased risk for type 2 diabetes.

Müssig K, Machicao F, Machann J, Schick F, Claussen CD, Stefan N, Fritsche A, Häring HU, Staiger H. BMC Med Genet. (2010)4;11:84 ([Artikel](#)) ([pdf](#))

Brief review: glucocorticoid excretion in obesity.

Müssig K, Remer T, Maser-Gluth C. J Steroid Biochem Mol Biol. (2010);121:589 ([Abstract PubMed](#))

[Diabetes, insulin, insulin analogues, and cancer].

[Article in German]

Müssig K, Staiger H, Kantartzis K, Fritsche A, Kanz L, Häring HU. Dtsch Med Wochenschr.(2010);135:924 ([Abstract PubMed](#))

Genetic variants affecting incretin sensitivity and incretin secretion.

Müssig K, Staiger H, Machicao F, Häring HU, Fritsche A. Diabetologia. (2010);53:2289 ([Artikel](#)) ([pdf](#))

Genetic variants in MTNR1B affecting insulin secretion.

Müssig K, Staiger H, Machicao F, Häring HU, Fritsche A. Ann Med. (2010);42:387 ([Abstract PubMed](#))

Phosphorylation of PRAS40 on Thr246 by PKB/AKT facilitates efficient phosphorylation of Ser183 by mTORC1.

Nascimento EB, Snel M, Guigas B, van der Zon GC, Kriek J, Maassen JA, Jazet IM, Diamant M, Ouwens DM. (2010) Cell Signal. 22: 961. ([Abstract PubMed](#))

Downregulation of mouse intestinal Na(+)-coupled glucose transporter SGLT1 by gum arabic (Acacia Senegal).

Nasir O, Artunc F, Wang K, Rexhepaj R, Föllner M, Ebrahim A, Kempe DS, Biswas R, Bhandaru M, Walter M, Mohebbi N, Wagner CA, Saeed AM, Lang F. Cell Physiol Biochem. (2010);25:203 ([Abstract PubMed](#))

Insulin sensitivity and beta-cell function in the offspring of type 2 diabetic patients: impact of line of inheritance.

Natali A, Muscelli E, Mari A, Balkau B, Walker M, Tura A, Anderwald C, Golay A, Ferrannini E; Relationship between Insulin Sensitivity and Cardiovascular disease investigators. (2010) J. Clin. Endocrinol. Metab. 95: 4703. ([Abstract PubMed](#))

Primäre Prävention des Typ-2-Diabetes - Patient und Therapeut - ihre wechselseitige Rolle in der Prävention

Neumann, P.E.H. Schwarz; Prävention und Gesundheitsförderung (2010);4:319

Body weight and energy homeostasis was not affected in C57BL/6 mice fed high whey protein or leucine-supplemented low-fat diets.

Noatsch A, Petzke KJ, Millrose MK, Klaus S. Eur J Nutr. (2010) [\(Artikel\)](#) [\(pdf\)](#)

Effects of acute psychological stress on glucose metabolism and subclinical inflammation in patients with post-traumatic stress disorder.

Nowotny B, Cavka M, Herder C, Loffler H, Poschen U, Joksimovic L, Kempf K, Krug AW, Koenig W, Martin S, Kruse J. (2010) Horm.Metab Res. 42: 746. [\(Abstract PubMed\)](#)

Quality indicators for the prevention of type 2 diabetes in Europe--IMAGE.

Pajunen P. et al.: Horm Metab Res. (2010) ;42 Suppl 1:S56 [\(Abstract PubMed\)](#)

A European evidence-based guideline for the prevention of type 2 diabetes.

Paulweber B. et al.: Horm Metab Res. (2010) ;42 Suppl 1:S3 [\(Abstract PubMed\)](#)

Socio-economic differences in life expectancy among persons with diabetes mellitus or myocardial infarction: results from the German MONICA/KORA study.

Perna L, Thien-Seitz U, Ladwig KH, Meisinger C, Mielck A. BMC Public Health. (2010) 10: 135. [\(Artikel\)](#) [\(pdf\)](#)

Genetic evidence that raised sex hormone binding globulin (SHBG) levels reduce the risk of type 2 diabetes.

Perry JR et al. (2010) Hum.Mol.Genet. 19: 535. [\(Artikel\)](#) [\(pdf\)](#)

Relationships of circulating sex hormone-binding globulin with metabolic traits in humans.

Peter A, Kantartzis K, Machann J, Schick F, Staiger H, Machicao F, Schleicher E, Fritsche A, Häring HU, Stefan N. (2010) Diabetes 59: 3167. [\(Abstract PubMed\)](#)

Dendritic cell-targeted pancreatic beta-cell antigen leads to conversion of self-reactive CD4(+) T cells into regulatory T cells and promotes immunotolerance in NOD mice.

Petzold C, Riewaldt J, Koenig T, Schallenberg S, Kretschmer K. (2010) Rev Diabet Stud. 7: 47. [\(Abstract PubMed\)](#)

Impact of age on the relationships of brown adipose tissue with sex and adiposity in humans.

Pfannenberger C, Werner MK, Ripkens S, Stef I, Deckert A, Schmadl M, Reimold M, Häring HU, Claussen CD, Stefan N. (2010) Diabetes 59: 1789. [\(Artikel\)](#) [\(pdf\)](#)

Role of the gut Peptide glucose-induced insulinomimetic Peptide in energy balance.

Pfeiffer AF, Rudovich N, Weickert MO, Isken F. Results Probl Cell Differ. (2010) ;52:183 [\(Abstract PubMed\)](#) [\(pdf\)](#)

Association of T-cell reactivity with beta-cell function in recent onset type 1 diabetes patients.

Pfleger C, Meierhoff G, Kolb H, Schloot NC. (2010) J. Autoimmun. 34: 127. [\(Abstract PubMed\)](#)

Exercise training increases mitochondrial content and ex vivo mitochondrial function similarly in patients with type 2 diabetes and in control individuals.

Phielix E, Meex R, Moonen-Kornips E, Hesselink MK, Schrauwen P. (2010) Diabetologia 53: 1714. ([Artikel](#)) ([pdf](#))

Commentary: Use of the body mass index to assess the risk of health outcomes: time to say goodbye?

Pischon T. Int J Epidemiol. (2010) ;39:528 ([Artikel](#)) ([pdf](#))

Body and liver fat mass rather than muscle mitochondrial function determine glucose metabolism in women with a history of gestational diabetes mellitus.

Prikoszovich T, Winzer C, Schmid AI, Szendroedi J, Chmelik M, Pacini G, Krssák M, Moser E, Funahashi T, Waldhäusl W, Kautzky-Willer A, Roden M. Diabetes Care. (2011);34:430 ([Artikel](#)) ([pdf](#))

Myristate is selectively incorporated into surfactant and decreases dipalmitoylphosphatidylcholine without functional impairment.

Pynn CJ, Picardi MV, Nicholson T, Wistuba D, Poets CF, Schleicher E, Perez-Gil J, Bernhard W. Am J Physiol Regul Integr Comp Physiol. (2010);299:R1306-16 ([Abstract PubMed](#))

Hyperglycemia impairs proteasome function by methylglyoxal.

Queisser MA, Yao D, Geisler S, Hammes HP, Lochnit G, Schleicher ED, Brownlee M, Preissner KT. Diabetes. (2010);59:670 ([Artikel](#)) ([pdf](#))

Development of a type 2 diabetes risk model from a panel of serum biomarkers from the Inter99 cohort: response to Kolberg et Al.

Rathmann W, Kowall B, Schulze MB. (2010) Diabetes Care 33: e28. ([Artikel](#))

Prediction models for incident Type 2 diabetes mellitus in the older population: KORA S4/F4 cohort study.

Rathmann W, Kowall B, Heier M, Herder C, Holle R, Thorand B, Strassburger K, Peters A, Wichmann HE, Giani G, Meisinger C. (2010) Diabet.Med. 27: 1116. ([Abstract PubMed](#))

Is primary aldosteronism associated with diabetes mellitus? Results of the German Conn's Registry.

Reincke M, Meisinger C, Holle R, Quinkler M, Hahner S, Beuschlein F, Bidlingmaier M, Seissler J, Endres S; Participants of the German Conn's Registry. Horm Metab Res. (2010) 42: 435. ([Abstract PubMed](#))

The natural protective mechanism against hyperglycemia in vascular endothelial cells: roles of the lipid peroxidation product 4-hydroxydodecadienal and peroxisome proliferator-activated receptor delta.

Riahi Y, Sin-Malia Y, Cohen G, Alpert E, Gruzman A, Eckel J, Staels B, Guichardant M, Sasson S. (2010) Diabetes 59: 808. (Artikel) ([pdf](#))

Automated suppression of sample-related artifacts in Fluorescence Correlation Spectroscopy.

Ries J, Bayer M, Csúcs G, Dirkx R, Solimena M, Ewers H, Schwille P. Opt Express. (2010) 24;18:11073
([Artikel](#)) ([pdf](#))

Reducing visceral adipose tissue mass is essential for improving endothelial function in type 2 diabetes prone individuals.

Rittig K, Hieronimus A, Thamer C, Machann J, Peter A, Stock J, Schick F, Fritsche A, Stefan N, Häring HU, Balletshofer B. Atherosclerosis. (2010);212:575 ([Abstract PubMed](#))

Effect of meglitinides on postprandial ghrelin secretion pattern in type 2 diabetes mellitus.

Rudovich N, Möhlig M, Otto B, Pivovarova O, Spranger J, Weickert MO, Pfeiffer AF. Diabetes Technol Ther. (2010) ;12:57 ([Artikel](#)) ([pdf](#))

Combination of acarbose and ezetimibe prevents non-alcoholic fatty liver disease: a break of intestinal insulin resistance?

Rudovich NN, Weickert MO, Machann J, Pfeiffer AF. J Hepatol. (2010);52:952 ([Abstract](#))

A high normal TSH is associated with the metabolic syndrome.

Ruhla S, Weickert MO, Arafat AM, Osterhoff M, Isken F, Spranger J, Schöfl C, Pfeiffer AF, Möhlig M. Clin Endocrinol (Oxf). (2010) ;72:696 ([Abstract PubMed](#))

Associations between BMI and the FTO gene are age dependent: results from the GINI and LISA birth cohort studies up to age 6 years.

Rzehak P, Scherag A, Grallert H, Sausenthaler S, Koletzko S, Bauer CP, Schaaf B, von Berg A, Berdel D, Borte M, Herbarth O, Krämer U, Illig T, Wichmann HE, Hebebrand J, Heinrich J; GINI and LISA Study Group. Obes Facts. (2010) 3: 173. ([Abstract PubMed](#))

Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge.

Saxena R, et al. (2010) Nature Genetics 42: 142. ([Artikel](#)) ([pdf](#))

Two new Loci for body-weight regulation identified in a joint analysis of genome-wide association studies for early-onset extreme obesity in French and German study groups.

Scherag A, et. al. PLoS Genet. (2010) 6 ([Artikel](#)) ([pdf](#))

Role of zinc finger transcription factor zfp69 in body fat storage and diabetes susceptibility of mice.

Scherneck S, Vogel H, Nestler M, Kluge R, Schürmann A, Joost HG. Results Probl Cell Differ. (2010);52:57 ([Abstract PubMed](#))

Impact of postprandial and fasting glucose concentrations on HbA1c in patients with type 2 diabetes.

Schernthaner G, Guerci B, Gallwitz B, Rose L, Nicolay C, Kraus P, Kazda C. Diabetes Metab. (2010);36:389 ([Abstract PubMed](#)) ([Artikel](#))

A race for RAGE ligands.

Schleicher ED. Kidney Int. (2010);78:234 ([Abstract PubMed](#))

Immunological intervention to preserve beta-cell function in type 1 diabetes.

Schlööt NC. (2010) Dtsch.Med.Wochenschr. 135: 915. ([Abstract Pubmed](#))

Diabetes Antibody Standardization Program: evaluation of assays for insulin autoantibodies.

Schlösser M, Müller PW, Törn C, Bonifacio E, Bingley PJ; Participating Laboratories. Diabetologia. (2010);53:2611. ([Artikel](#)) ([pdf](#))

Diabetesstudien im Überblick - Intensive Blutzuckersenkung von Vorteil?

Schnoor Luitgard, Schwarz Peter E.H. Diabetes aktuell (2010);8:22

Liver ATP synthesis is lower and relates to insulin sensitivity in patients with type 2 diabetes mellitus.

Schmid A, Szendroedi J, Chemlik M, Krssak M, Moser E, Roden M. (2010) Diabetes Care 34: 448. ([Artikel](#)) ([pdf](#))

β2-Syntrophin is a Cdk5 substrate that restrains the motility of insulin secretory granules.

Schubert S, Knoch KP, Ouwendijk J, Mohammed S, Bodrov Y, Jäger M, Altkrüger A, Wegbrod C, Adams ME, Kim Y, Froehner SC, Jensen ON, Kalaidzidis Y, Solimena M. (2010) PLoS One.5: e12929. ([Artikel](#)) ([pdf](#))

The effect of 6-weeks of aerobic exercise training on serum fetuin-A levels in non-diabetic obese women.

Schultes B, Frick J, Ernst B, Stefan N, Fritsche A. Exp Clin Endocrinol Diabetes. (2010);118:754 ([Abstract PubMed](#))

Diabetesprävalenz: Verlässliche Schätzungen stehen noch aus.

Schulze M.B., Rathmann W., Giani G., Joost H.-G. Deutsches Ärzteblatt 107;A1694

Fasting plasma glucose and Type 2 diabetes risk: a non-linear relationship.

Schulze MB, Fritsche A, Boeing H, Joost HG. (2010) Diabet Med. 27: 473. ([Abstract PubMed](#))

Early Detection of Diabetes - Screening Instruments and Risk Scores

Schwarz, Peter E.H. Diabetes aktuell (2010);08:164

Intervention for the Prevention of Diabetes Mellitus - The Prevention Manager

Schwarz, Peter E.H.; Diabetes aktuell (2010);8:169

6th World Congress on Prevention of Diabetes Mellitus and its Complications Peter

Schwarz, Peter E.H. Expert Review of Endocrinology & Metabolism (2010);5:4:517

Scientific summary of the 6th World Congress on Prevention of Diabetes and its Complications (WCPD 2010)

Schwarz Peter E.H. Anatol J Obstet Gynecol (2010);2:1

Diabetes: 10 years of incidence rates related

Schwarz Peter E.H. Diabetologie und Stoffwechsel (2010);5,3,p157

Intervention for the Prevention of Diabetes mellitus - Lifestyle intervention

Schwarz Peter E.H. Diabetes aktuell (2010); 8:172

EZSCAN - eine neue, einfache, validierte Methode zur Diagnostik eines Diabetesrisikos

Schwarz Peter E.H., P.M.Röttger; Diabetes Aktuell (2010);8:208

Entwicklung zum Diabetes lässt sich abwenden

Schwarz Peter E.H. Forschung & Praxis (2010);506:11

Prävention des Diabetes - wie können wir Präventionsmaßnahmen in Praxis umsetzen

Schwarz Peter E.H. Adipositas (2010);4:95

FINDRISK - ein praktisches Screeninginstrument für Diabetesrisiko

Schwarz Peter E.H. klinikarzt (2010);39:546

From evidence to practice - the IMAGE project-new standards in the prevention of type 2 diabetes

Schwarz PE, Lindström J. Diabetes Res Clin Pract.(2010);29

Highlights of the World Congress for Diabetes Prevention - From Theory to Practical Realisation

Schwarz Peter E.H. Diabetes aktuell (2010);8:158

Towards risk stratification in systemic atherosclerosis: value of myocardial function and viability imaging as an adjunct to MR angiography.

Seeger A, Fenchel M, Kramer U, Bretschneider C, Grimm F, Klumpp B, Scheule A, Balletshofer B, Claussen CD, Miller S. Eur Radiol. (2010);20:838 ([Artikel](#)) ([pdf](#))

Chemerin correlates with markers for fatty liver in morbidly obese patients and strongly decreases after weight loss induced by bariatric surgery.

Sell H, Divoux A, Poitou C, Basdevant A, Bouillot JL, Bedossa P, Tordjman J, Eckel J, Clement K. (2010) J.Clin.Endocrinol.Metab 95: 2892. ([Abstract PubMed](#))

The associations of cholesterol metabolism and plasma plant sterols with all-cause and cardiovascular mortality.

Silbernagel G, Fauler G, Hoffmann MM, Lütjohann D, Winkelmann BR, Boehm BO, März W. J Lipid Res. (2010);51:2384 ([Abstract PubMed](#))

Evaluating the maintenance of effect of duloxetine in patients with diabetic peripheral neuropathic pain.

Skljarevski V, Desai D, Zhang Q, Chapell A, Detke M, Gross J, Ziegler D. (2009) Diabetes Metab Res Rev 25: 623. ([Abstract PubMed](#))

Insulin release: shedding light on a complex matter.

Solimena M, Speier S. (2010) Cell Metab. 12: 5. ([Abstract PubMed](#))

Common variants at ten genomic loci influence hemoglobin A1C levels via glycemic and non-glycemic pathways.

Soranzo N, et al. (2010) Diabetes 59: 3229. ([Abstract PubMed](#))

Mesenchymal cells appearing in pancreatic tissue culture are bone marrow-derived stem cells with the capacity to improve transplanted islet function.

Sordi V, Melzi R, Mercalli A, Formicola R, Doglioni C, Tiboni F, Ferrari G, Nano R, Chwalek K, Lammert E, Bonifacio E, Borg D, Piemonti L. Stem Cells. (2010) ;28:140. ([Artikel](#)) ([pdf](#))

Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index.

Speliotes EK. et al.: Nat Genet. (2010) ;42:937. ([Artikel](#)) ([pdf](#))

Imaging of primary and secondary inflammatory diseases involving large and medium-sized vessels and their potential mimics: a multitechnique approach.

Spira D, Kötter I, Ernemann U, Balletshofer B, Pfannenberg CA, Fenchel M, Horger M. AJR Am J Roentgenol. (2010);194:848 ([Artikel](#)) ([pdf](#))

Quantitative assessment of intrahepatic lipids using fat-selective imaging with spectral-spatial excitation and in-/opposed-phase gradient echo imaging techniques within a study population of extremely obese patients: feasibility on a short, wide-bore MR scanner.

Springer F, Machann J, Schwenzler NF, Ballweg V, Würslin C, Schneider JH, Fritsche A, Claussen CD, Schick F. Invest Radiol. (2010);45:484 ([Abstract PubMed](#))

Quantitative assessment of visceral fat in morbidly obese patients by means of wide-bore MRI and its relation to lower esophageal sphincter pressure and signs of gastroesophageal reflux.

Springer F, Schwarz M, Machann J, Fritsche A, Claussen CD, Schick F, Schneider JH. Obes Surg. (2010);20:749 ([Artikel](#)) ([pdf](#))

Association of lower plasma fetuin-a levels with peripheral arterial disease in type 2 diabetes: response to Eraso et al.

Stefan N, Weikert C, Ix JH, Fritsche A, Häring HU. Diabetes Care. (2010) ;33 ([Artikel](#)) ([pdf](#))

Circulating palmitoleate strongly and independently predicts insulin sensitivity in humans.

Stefan N, Kantartzis K, Celebi N, Staiger H, Machann J, Schick F, Cegan A, Elcnerova M, Schleicher E, Fritsche A, Häring HU. Diabetes Care. (2010);33:405 ([Artikel](#)) ([pdf](#))

Letter by Stefan et al regarding article, "Impact of body mass index and the metabolic syndrome on the risk of cardiovascular disease and death in middle-aged men".

Stefan N, Fritsche A, Häring HU. Circulation. (2010)14;122:e456 ([Artikel](#)) ([pdf](#))

The temporal sequence of magnetic brain activity for food categorization and memorization--an exploratory study.

Stingl KT, Rogić M, Stingl K, Canova C, Tschritter O, Braun C, Fritsche A, Preissl H. (2010) Neuroimage 52: 1584. ([Abstract PubMed](#))

The temporal sequence of magnetic brain activity for food categorization and memorization--an exploratory study.

Stingl KT, Rogić M, Stingl K, Canova C, Tschritter O, Braun C, Fritsche A, Preissl H. Neuroimage. (2010)1;52:1584 ([Abstract PubMed](#))

The insulin secretory granule as a signaling hub.

Suckale J, Solimena M. Trends Endocrinol Metab. (2010) ;21:599 ([Abstract PubMed](#))

Combined risk allele score of eight type 2 diabetes genes is associated with reduced first-phase glucose-stimulated insulin secretion during hyperglycemic clamps.

't Hart LM, Simonis-Bik AM, Nijpels G, van Haeften TW, Schäfer SA, Houwing-Duistermaat JJ, Boomsma DI, Groenewoud MJ, Reiling E, van Hove EC, Diamant M, Kramer MH, Heine RJ, Maassen JA, Kirchhoff K, Machicao F, Häring HU, Slagboom PE, Willemsen G, Eekhoff EM, de Geus EJ, Dekker JM, Fritsche A. Diabetes. (2010);59:287 ([Artikel](#)) ([pdf](#))

Large effects on body mass index and insulin resistance of fat mass and obesity associated gene (FTO) variants in patients with polycystic ovary syndrome (PCOS).

Tan S, Scherag A, Janssen OE, Hahn S, Lahner H, Dietz T, Scherag S, Grallert H, Vogel CI, Kimmig R, Illig T, Mann K, Hebebrand J, Hinney A. BMC Med Genet. (2010) 11:12 ([Artikel](#)) ([pdf](#))

Interscapular fat is strongly associated with insulin resistance.

Thamer C, Machann J, Staiger H, Müssig K, Schwenzer N, Ludescher B, Machicao F, Claussen C, Fritsche A, Schick F, Stefan N, Häring HU. (2010) J Clin Endocrinol Metab. 95: 4736. ([Abstract PubMed](#))

Associations between leptin and the leptin / adiponectin ratio and incident Type 2 diabetes in middle-aged men and women: results from the MONICA / KORA Augsburg study 1984-2002.

Thorand B, Zierer A, Baumert J, Meisinger C, Herder C, Koenig W. (2010) Diabet.Med. 27: 1004. ([Abstract PubMed](#))

Loss of the actin remodeler Eps8 causes intestinal defects and improved metabolic status in mice.

Tocchetti A, Soppo CB, Zani F, Bianchi F, Gagliani MC, Pozzi B, Rozman J, Elvert R, Ehrhardt N, Rathkolb B, Moerth C, Horsch M, Fuchs H, Gailus-Durner V, Beckers J, Klingenspor M, Wolf E, Hrabě de Angelis M, Scanziani E, Tacchetti C, Scita G, Di Fiore PP, Offenhäuser N. PLoS One. (2010) 5: e9468. ([Artikel](#)) ([pdf](#))

Primary prevention of type 2 diabetes is advancing towards the mature stage in Europe.

Tuomilehto J, Schwarz PE. Horm Metab Res. (2010) ;42 Suppl 1:S1. ([Artikel](#)) ([pdf](#))

Association of serum levels of retinol-binding protein 4 with male sex but not with insulin resistance in obese patients.

Ulgen F, Herder C, Kuhn MC, Willenberg HS, Schott M, Scherbaum WA, Schinner S. (2010) Arch.Physiol Biochem. 116: 57. ([Abstract PubMed](#))

Transmission infrared spectroscopy of whole blood--complications for quantitative analysis from leucocyte adhesion during continuous monitoring.

Vahlsing T, Damm U, Kondepoti VR, Leonhardt S, Brendel MD, Wood BR, Heise HM. J Biophotonics. (2010) ;3:567. ([Abstract PubMed](#))

Phenotypic and pathomorphological characteristics of a novel mutant mouse model for maturity-onset diabetes of the young type 2 (MODY 2).

van Bürck L, Blutke A, Kautz S, Rathkolb B, Klaften M, Wagner S, Kemter E, Hrabé de Angelis M, Wolf E, Aigner B, Wanke R, Herbach N. *Am J Physiol Endocrinol Metab.* (2010) 298:E512 ([Artikel](#)) ([pdf](#))

High levels of dietary stearate promote adiposity and deteriorate hepatic insulin sensitivity.

van den Berg SA, Guigas B, Bijland S, Ouwens M, Voshol PJ, Frants RR, Havekes LM, Romijn JA, van Dijk KW. (2010) *Nutr.Metab (Lond)* 7: 24. ([Artikel](#)) ([pdf](#))

Diabetic cardiomyopathy in Zucker diabetic fatty rats: the forgotten right ventricle.

van den Brom CE, Bosmans JW, Vlasblom R, Handoko LM, Huisman MC, Lubberink M, Molthoff CF, Lammertsma AA, Ouwens MD, Diamant M, Boer C. (2010) *Cardiovasc. Diabetol.* 9: 25. ([Artikel](#)) ([pdf](#))

Cyclooxygenase-2 controls energy homeostasis in mice by de novo recruitment of brown adipocytes.

Vegiopoulos A, Müller-Decker K, Strzoda D, Schmitt I, Chichelnitskiy E, Ostertag A, Berriel Diaz M, Rozman J, Hrabé de Angelis M, Nüsing RM, Meyer CW, Wahli W, Klingenspor M, Herzig S. *Science.* (2010); 328:1158 ([Artikel](#)) ([pdf](#))

Diabetic polyneuropathy. The crystal ball of cardiovascular risk.

Vinik A, Maser R, Ziegler D. (2010) *Diabetes Care* 33: 1688. ([Artikel](#)) ([pdf](#))

Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis.

Voight BF, et al. (2010) *Nature Genetics* 42: 579. ([Abstract PubMed](#))

Circulating vaspin is unrelated to insulin sensitivity in a cohort of nondiabetic humans.

von Loeffelholz C, Möhlig M, Arafat AM, Isken F, Spranger J, Mai K, Randevara HS, Pfeiffer AF, Weickert MO. *Eur J Endocrinol.* (2010);162:507 ([Artikel](#)) ([pdf](#))

No effect of the 1 α ,25-dihydroxyvitamin D₃ on beta-cell residual function and insulin requirement in adults with new-onset type 1 diabetes.

Walter M, Kaupper T, Adler K, Foersch J, Bonifacio E, Ziegler AG. *Diabetes Care.* (2010) ;33:1443. ([Artikel](#)) ([pdf](#))

Vascular function in obese children with non-alcoholic fatty liver.

Weghuber D, Roden M, Franz C, Chmelik M, Torabia S, Nowotny P, Gruber S, Waldhäusl W, Klingler A, Biegelmayer C, Bischof S, Wolzt M, Schaller G, Widhalm K. (2010) *J. Pediatr. Obes.* Epub Oct 2010. ([Abstract PubMed](#))

BMI at age 8 years is influenced by the type 2 diabetes susceptibility genes HHEX-IDE and CDKAL1.

Winkler C, Bonifacio E, Grallert H, Henneberger L, Illig T, Ziegler AG. *Diabetes.* (2010);59:2063. ([Artikel](#)) ([pdf](#))

Presence of gallstones or kidney stones and risk of type 2 diabetes.

Weikert C, Weikert S, Schulze MB, Pischon T, Fritsche A, Bergmann MM, Willich SN, Boeing H. Am J Epidemiol. (2010) 15;171:447 ([Artikel](#)) ([pdf](#))

The D299G/T399I Toll-like receptor 4 variant associates with body and liver fat: results from the TULIP and METSIM Studies.

Weyrich P, Staiger H, Stančáková A, Machicao F, Machann J, Schick F, Stefan N, Kuusisto J, Laakso M, Schäfer S, Fritsche A, Häring HU. PLoS One. (2010)15;5:e13980 ([Artikel](#)) ([pdf](#))

Study on pancreatic islet adaptation and gene expression during pregnancy in rats.

Xue Y, Liu C, Xu Y, Yuan Q, Xu K, Mao X, Chen G, Wu X, Brendel MD, Liu C. Endocrine. (2010);37:83 ([Artikel](#)) ([pdf](#))

A new collagenase blend increases the number of islets isolated from mouse pancreas.

Yesil P, Michel M, Chwalek K, Pedack S, Jany C, Ludwig B, Bornstein SR, Lammert E. Islets. (2009) ;1:185. ([Artikel](#)) ([pdf](#))

Metabonomic fingerprints of fasting plasma and spot urine reveal human pre-diabetic metabolic traits.

Zhao X, Fritsche J, Wang J, Chen J, Rittig K, Schmitt-Kopplin P, Fritsche A, Häring HU, Schleicher ED, Xu G, Lehmann R. Metabolomics. (2010);6:362 ([Artikel](#)) ([pdf](#))

Efficacy and safety of lacosamide in painful diabetic neuropathy.

Ziegler D, Hidvégi T, Gurieva I, Bongardt S, Freyhagen R, Sen D, Sommerville K. (2010) Diabetes Care 33: 839. ([Artikel](#)) ([pdf](#))

Genetic variation of the FADS1 FADS2 gene cluster and n-6 PUFA composition in erythrocyte membranes in the European Prospective Investigation into Cancer and Nutrition-Potsdam study.

Zietemann V, Kröger J, Enzenbach C, Jansen E, Fritsche A, Weikert C, Boeing H, Schulze MB. Br J Nutr. (2010);104:1748 ([Abstract PubMed](#))